

HYGROTECH

SUSTAINABLE SOLUTIONS

TOMATOES

TOP VARIETIES

PAMELA

- * Strong indeterminate growth with dense foliage
- * Good tolerance to diseases
- * Cylindrical in shape
- * Extremely good taste
- * Good for fresh market

F350


- * Determinate saladette
- * BW tolerant
- * Resistant to: Verticillium wilt race 1, root knot nematodes,
- * YMV, TSWV, Bacterial speck, TYLCV

JULIA

- * Determinate saladette
- * BW tolerant
- * Resistant to: Verticillium wilt race 1, root knot nematodes,
- * YMV, TSWV, Bacterial speck, TYLCV

SALADITA CHERRY TOMATO

- * Vigorous, open, medium generative growth habit
- * Exceptionally deep red colour 15-25g
- * No green shoulders
- * Firm, oval shaped fruit



PLEASE CONTACT US AND GET ADVISED FROM OUR OFFICES

HEAD OFFICE

Tigoni Centre, Limuru Road
P. O. Box 41446, Nairobi, Kenya
Tel: +254 20 2053916/7/8, 20 2053921
Cell: + 254 722 205148/+ 254 728 607582
Email: info@hygrotech.co.ke

NAIVASHA BRANCH






P. O. Box 1484-20117
Naivasha, Kenya
Tel: + 254 50 50500
Cell: + 254 728 607583/578
Email: info@hygrotech.co.ke

TRIACHEM (TZ) BRANCH

P. O. Box 2750
Arusha, Tanzania
Tel: + 255 784 695193
Cell: + 255 767 869869
Email: info@triachem.com

GENERAL TOMATO GROWING GUIDE

FOLIAR FEEDING		TOMATO DISEASES		REMEDY
Sowing	Transplanting	Seedling	Fusarium, Pythium Rhizoctonia, Phytophthora	Thiamethoxam Sporekill Fosfaspray Kic start
1st two weeks	Kicstart - 2ml/L water Fosfaspray - 5g/L water Sporekill - 0.5ml/L Water	Foliage diseases		Sporekill Liquicop Hygroboost Flo
1st Month	Nitrospray - 2ml/L water Hyperfeed - 5g/L water Hygroboost Flo - 2ml/L Water Seamaxx - 2.5ml/L water	Wilt virus		
		Mosaic virus		
		Bacterial canker		
2nd Month	Hyperfeed - 5g/L water Hygroboost Flo - 2ml/L Water Nitrospray - 2ml/L water Sporekill - 1ml/L water Liquicop - 2ml/L water	Early & late blights (Fungi)		
		Soil borne diseases		
		Fusarium/pythium/ Ralstonia	Perika Mustard caliente	
3rd and 4th Month	Sporekill - 1ml/L water Hyperfeed - 5g/L water Calmabon - 2ml/L water Coprtrace - 2ml/1L water	Root knot Nematodes (<i>M. spp</i>)		
		Post-harvest		
		Black and grey mould	Vapour guard	
		General Pests		Acaricides Chlorantraniliprole
		Thrips and aphids		
		White flies		
		Cutworms		
		Leaf miners (<i>Lyriomyza spp</i>) Hornworms		
		Physiological disorder		Calcium / Calmabon
		Blossom-end-rot		

<p>Shape / umbo</p>  <p>Cylindrical-oval</p>	<p>Days to harvest / siku za kukomaa</p> <p>80-120 days</p>  <p>Transplanting / Hupandwa niche</p>	<p>Sowing Season / msimu wa kupanda</p> 
<p>Depth / undani wa kupanda</p>  <p>2 cm</p>	<p>Spacing / umbali mmea kwa mmea</p> <p>40x60 cm</p> 	<p>Watering / kuweka maji</p> <p>Water regularly / mara kwa mara</p> 