

GENERAL TOMATO GROWING GUIDE

FOLIAR FEEDING	
Sowing	Transplanting
1 st two weeks	Kic start – 2ml/L water Fosfaspray – 5g/L water Sporekill – 0.5ml/L water
1 st Month	Nitrospray – 2ml/L water Hyperfeed – 5g/L water Hygroboost Flo– 2ml/L water Seamaxx – 2.5ml/L water
2 nd Month	Hyperfeed – 5g/L water Hygroboost Flo– 2ml/L water Nitrospray – 2ml/L water Sporekill –1ml/L water Liquicop –2ml/1L water
3 rd and 4 th Month	Sporekill –1ml/L water Hyperfeed – 5g/L water Calmabon- 2ml/L water Copttrace - 2ml/ 1L water

TOMATO DISEASES		REMEDY
Seedling	Fusarium, Pythium Rhizoctonia, Phytophthora	Thiamethoxam Sporekill Fosfaspray Kic start
	Foliage diseases	
	Wilt virus	Sporekill Liquicop Hygroboost Flo
	Mosaic virus	
	Bacterial canker	
	Early & late blights (Fungi)	
Soil borne diseases		
	Fusarium/pythium/ Ralstonia	Perlka Mustard caliente
	Root knot Nematodes (<i>M. spp</i>)	
Post-harvest		
	Black and grey mould	Vapour guard Avoshine
General Pests		
	Thrips and aphids	Acaricides Chlorantraniliprole
	White flies	
	Cutworms	
	Leaf miners (<i>Lyriomyza spp</i>)	
	Hornworms	
Physiological disorder		
	Blossom-end-rot	Calcium/Calmabon

Shape /umbo

Cylindrical-oval



Days to harvest / siku za kukomaa

80-120 days



Transplanting / Hupandwa miche

Sowing Season / msimu wa kupanda



Depth / undani wa kupanda



2 cm

Spacing / umbali mmea kwa mmea

40x60 cm



Watering / kuweka maji

Water regularly / mara kwa mara

